

Chromosomal Mapping of Canine-Derived BAC Clones to the Red Fox and American Mink Genomes

ANNA V. KUKKOVA, NADEGDA V. VOROBIEVA, VIOLETTA R. BEKLEMISHEVA, JENNIFER L. JOHNSON, SVETLANA V. TEMNYKH, DMITRY V. YUDKIN, LYUDMILA N. TRUT, CATHERINE ANDRE, FRANCIS GALIBERT, GUSTAVO D. AGUIRRE, GREGORY M. ACLAND, AND ALEXANDER S. GRAPHODATSKY

From the James A. Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853 (Kukekova, Johnson, Temnykh, and Acland); the Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia (Vorobieva, Beklemisheva, Yudkin, Trut, and Graphodatsky); Une unités mixtes de recherche 6061, Centre national de la recherche scientifique, Institute of Genetics and Development, Faculty of Medicine, Rennes, France (Andre and Galibert); and the School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA (Aguirre).

Address correspondence to Anna V. Kukekova at the address above, or e-mail: avk5@cornell.edu.

Abstract

High-quality sequencing of the dog (*Canis lupus familiaris*) genome has enabled enormous progress in genetic mapping of canine phenotypic variation. The red fox (*Vulpes vulpes*), another canid species, also exhibits a wide range of variation in coat color, morphology, and behavior. Although the fox genome has not yet been sequenced, canine genomic resources have been used to construct a meiotic linkage map of the red fox genome and begin genetic mapping in foxes. However, a more detailed gene-specific comparative map between the dog and fox genomes is required to establish gene order within homologous regions of dog and fox chromosomes and to refine breakpoints between homologous chromosomes of the 2 species. In the current study, we tested whether canine-derived gene-containing bacterial artificial chromosome (BAC) clones can be routinely used to build a gene-specific map of the red fox genome. Forty canine BAC clones were mapped to the red fox genome by fluorescence in situ hybridization (FISH). Each clone was uniquely assigned to a single fox chromosome, and the locations of 38 clones agreed with cytogenetic predictions. These results clearly demonstrate the utility of FISH mapping for construction of a whole-genome gene-specific map of the red fox. The further possibility of using canine BAC clones to map genes in the American mink (*Mustela vison*) genome was also explored. Much lower success was obtained for this more distantly related farm-bred species, although a few BAC clones were mapped to the predicted chromosomal locations.

Key words: *Canis lupus familiaris*, comparative genomics, FISH, *Mustela vison*, *Vulpes vulpes*

The phenotypic variation observed in domestic and farm-bred fur animals is striking, but its genetic basis is far from being well understood. New genetic tools and genome sequences (<http://www.broad.mit.edu/mammals/>) create an opportunity for the analysis of monogenic and complex phenotypes in dog, cat, and livestock species (reviewed in Georges 2007; Karlsson et al. 2007; Pontius et al. 2007; Wayne and Ostrander 2007). Furthermore, information from well-studied genomes can be applied for comparative analysis and genetic mapping of traits of interest in closely related species (de Gortari et al. 1998; Rogers et al. 2000, 2006; Reed et al. 2005; Anistoroaei et al. 2007). Here, we explore an opportunity to use canine genomic tools for assignment of gene-specific markers in the genomes of 2 farm-bred species, the red fox (*Vulpes vulpes*) and American

mink (*Mustela vison*), in which artificial selection has produced a variety of coat color, morphological, and behavioral phenotypes (Shackelford 1948, 1949; Belyaev et al. 1975; Robinson 1975; Nes et al. 1988; Våge et al. 1997; Trut 1999, 2001; Kharlamova et al. 2000, 2007; Trapezov et al. 2008).

Canids (dog and fox) and mustelids (mink) belong to the same Canoidea branch of the Carnivora and diverged from a common ancestor about 40 million years ago (Ma), whereas red fox and dog, members of the Canidae family, diverged about 10 Ma (Wayne et al. 1997; Bininda-Emonds et al. 1999). The dog karyotype (*Canis lupus familiaris*) comprises 37 pairs of acrocentric autosomes and a pair of sex chromosomes, whereas the red fox has 16 pairs of metacentric autosomes, a pair of sex chromosomes, and 0–8

Table 1. FISH localization of canine gene-containing BAC clones in the genomes of the silver fox and American mink

Gene name	BAC clone	Fox FISH mapping (VVU)	Mink FISH mapping (MVI)	CanFam2 canine ortholog location	Dog chromosome based on RH mapping data (CFA)	Closest RH markers	θ	LOD	Marker location on canine FISH-RH-integrated map, cR (Breen et al. 2004)	Reference	Selected candidate genes	Genes that were previously mapped in the red fox by FISH
T-brachyury protein (<i>T</i> gene)	90-I8	1p1.5–2.6		Chr 1: 57, 228, 892-57, 237, 728	CFA1	FH3993	0.09	17.81	1164	AJ245513; Haworth et al. (2001)		
<i>HTR1B</i>	18-L8	1q3.5–3.6	1p26–27	Chr 12: 41, 662, 262-41, 662, 720	CFA12 (van den Berg et al. 2004)	FH3284 BAC_372-C2	0.12 0.02	16.75 25.28	1164 699	van den Berg et al. (2004)	n	
<i>CRHR1</i>	189-B1	2p2.10		Chr 9: 12, 791, 000-12, 841, 243	CFA9	EST5H8 BAC_372-C21	0.02 0.26	25.28 11.64	694 698	This study	n	
<i>GRIN2C</i>	429-O8	2p2.10		Chr 9: 8, 582, 559-8, 594, 291	CFA9	CCL2 GALK1	0.28 0.26	10.93 11.44	690	Sidjanin et al. (2003)		
<i>Tbx2</i> and <i>Tbx4</i>	327-P20	2p2.3–2.7		TBX2—Chr 9: 38, 559, 410-38, 567, 853	CFA9	SRP68 FH2846	0.37 0.07	8.09 22.01	645	AY192796, AY192797; Andelfinger et al. (2003)		
<i>SLC6A4</i>	59-N4	2p1.5–2.5		TBX4—Chr 9: 38, 487, 909-38, 513, 488 Chr 9: 47, 549, 263-47, 572, 016	CFA9	EST9A7 EST27D2	0.10 0.03	19.98 19.82	626 820	This study	n	
<i>SLC6A3</i>	89-K1	2p1.5–2.5		Chr 34: 14, 216, 551-14, 251, 928; Chr 2: 63, 201, 559-63, 242, 406 (CFA2 less likely)	CFA22	FH3235 REN128H16	0.12 0.29	16.90 11.65	811 105	This study	n	
<i>DBH</i>	73-H9	2p1.5–2.5		Chr 9: 53, 342, 444-53, 361, 759	CFA9	AH2H211 LEI-2D2	0.35 0.19	10.04 11.80	128 967	This study	n	
<i>C-KIT</i>	118-O18	2p1.1		Chr 13: 50, 040, 774-50, 124, 361	CFA13	REN177B24 REN126A23 KDR	0.22 0.07 0.08	11.06 16.21 15.42	967 704 711	This study	c	Graphodatsky et al. (2005)

Table 1. Continued

Gene name	BAC clone	Fox FISH mapping (VVU)	Mink FISH mapping (MVI)	CanFam2 canine ortholog location	Dog chromosome based on RH mapping data (CFA)	Closest RH markers	θ	LOD	Marker location on canine FISH-RH-integrated map, cR (Breen et al. 2004)	Reference	Selected candidate genes	Genes that were previously mapped in the red fox by FISH
<i>KDR</i>	67-P13	2p1.2		Chr 13: 50, 374, 957-50, 416, 670	CFA13 (Breen et al. 2004)					Graphodatsky et al. (2005)		Graphodatsky et al. (2005)
<i>PDGFRA</i>	311-A23	2p1.2		Chr 13: 49, 684, 632-49, 705, 871	CFA13	REN126A23	0.07	17.04	704	Graphodatsky et al. (2005)	c	Graphodatsky et al. (2005)
<i>SLC6A2</i>	289-M11	2q3.4-3.7		Chr 2: 63, 198, 411-63, 242, 614	CFA2	C13.365 FH2608	0.07 0.03	16.27 19.16	692 841	AC129099	n	
<i>GAD1</i>	130-I2	3p2.2-2.6		Chr 36: 18, 369, 306-18, 412, 283	CFA36	BAC_372-A19 EST19A10	0.07 0.08	16.98 23.79	841 348	This study	n	
<i>CYP3A26</i> (<i>CYP3A4</i>)	168-J21	3q1.5		Chr 6: 12, 784, 228-12, 821, 910	CFA6	BAC_381-A3 BAC_382-E21	0.17 0.21	18.82 16.02	330 308	AC131027		
<i>ABAT</i>	65-N8	3q2.1-3.1		Chr 6: 36, 426, 512-36, 474, 610	CFA6	BAC_372-O10 REN88M24	0.22 0.09	15.53 13.65	317 709	This study	n	
<i>CDH23</i>	99-C20	4q1.6-2.2		Chr 4: 25, 492, 895-25, 781, 226	CFA4 (Kuiper et al. 2002)	REN84K01 BAC_99-C20	0.09 0.31	12.81 9.73	689 552	Kuiper et al. (2002)		
<i>DRD1</i>	1-M13	4q1.3-1.7		Chr 4: 40, 743, 436-40, 746, 624	CFA4	BAC_381-H2 FH3106	0.35 0.04	8.71 18.39	532 874	This study	n	
<i>ATOX1</i> pseudogene	84-B18	4q2.1-2.2	1q31-33	Chr 4: 60, 803, 406-60, 812, 281	CFA4	FH2412 BAC_374-E21	0.11 0.05	15.16 14.10	880 1547	AF286877; van de Sluis et al. (2001)		
<i>GABRG2</i>	49-K24	4q3.1-3.3		Plus CFA18, 19, 20, and 24 Chr 4: 51, 998, 352-52, 105, 402	CFA4	FH1015	0.05	14.10	1547			
<i>GABRG2</i>						EST11E12	0.08	15.48	1449	This study	n	
<i>HTR2A</i>	127-D1	6p2.2-2.5		Chr 22: 7, 395, 321-7, 452, 531	CFA22	REN282C02 REN42F10	0.11 0.06	14.39 26.49	1476 114	This study	n	Klukowska-Rötzler et al. (2005)
						REN128H16	0.04	26.25	105			

Table 1. Continued

Gene name	BAC clone	Fox FISH mapping (VVU)	Mink FISH mapping (MVI)	CanFam2 canine ortholog location	Dog chromosome based on RH mapping data (CFA)	Closest RH markers	θ	LOD	Marker location on canine FISH-RH-integrated map, cR (Breen et al. 2004)	Reference	Selected candidate genes	Genes that were previously mapped in the red fox by FISH
<i>TYRP2 (DCT)</i>	65-H10	6p1.3–1.4		Chr 22: 48, 220, 073-48, 253, 910	CFA22	DCT	0.03	20.52	592	This study	c	
<i>HOXA</i>	78-A3	7p2.3				FH2662	0.09	17.97	578	Li et al. (1999)		
<i>GNGT1</i>		7q1.1–1.3	3q14	Chr 14: 22, 363, 018-22, 367, 549	CFA14	BAC_372-I9	0.06	20.30	450	This study		
<i>SHARP1 (BHLHB3)</i>	115-A9	8p1.2–2.1		Chr 27: 24, 475, 788-24, 480, 605	CFA27	BAC_375-O13 EST7A1	0.06 0.28	20.03 9.64	494 394	Kukekova et al. (2003)		
<i>ST8SLA1</i>	333-F7	8p1.1–1.2		Chr 27: 27, 948, 915-28, 038, 424	CFA27	PTHLH STS156J06	0.34 0.12	7.73 18.72	379 492	This study		
<i>UGT1A6</i>	252-I15	9p2.3–2.5		Chr 25: 48, 138, 801-48, 183, 812	CFA25	D04813 ALPI	0.14 0.06	17.65 18.49	502 835	AC093453		
<i>SLC6A1</i>	102-D19	9q1.2–p1.1		Chr 20: 10, 344, 464-10, 386, 539	CFA20	FH4027 REN55P21	0.07 0.21	17.64 17.12	844 226	This study	n	
<i>IGF1</i>	126-J3	10q2.1–2.2		Chr 15: 44, 229, 347-44, 285, 955	CFA15	EST13G6 FH2295	0.22 0.09	16.39 17.74	263 795	This study	m	Szczerbal et al. (2003)
<i>SLC25A3</i>	50-E1	10q1–2.3		Chr 15: 41, 002, 937-41, 009, 210	CFA15	IGF1 REN134K05	0.14 0.09	15.70 21.03	808 736	Debenham et al. (2001)		
<i>TPH1</i>	147-H22	11p2.3–2.4		Chr 21: 43, 739, 337-43, 757, 824	CFA21	REN143N23 SAA2	0.10 0.13	18.76 19.56	711 825	This study	n	
<i>TECTA</i>	59-C2	12q2.1–2.5		Chr 5: 15, 886, 369-15, 954, 803	CFA5 (Drögemüller et al. 2002)	EST11A11 REN42N13	0.14 0.06	18.58 19.67	783 385	Drögemüller et al. (2002)		
<i>MYO15A</i>	362-O13	12q2.9	8p11	Chr 5: 44, 369, 188-44, 426, 838	CFA5 (Rak et al. 2002)	REN78M01 BAC_375-M3	0.06 0.05	19.17 13.05	397 933	Rak et al. (2002)		

Table 1. Continued

Gene name	BAC clone	Fox FISH mapping (VVU)	Mink FISH mapping (MVI)	CanFam2 canine ortholog location	Dog chromosome based on RH mapping data (CFA)	Closest RH markers	θ	LOD	Marker location on canine FISH-RH-integrated map, cR (Breen et al. 2004)	Reference	Selected candidate genes	Genes that were previously mapped in the red fox by FISH
<i>MC1R</i>	130-L5	12q3.1–3.3		Chr 5: 66, 692, 398–66, 693, 344	CFA5	H006 MC1R	0.15 0.07	10.98 16.21	933 1421	This study	c	
<i>CNGB3</i>	100-O2	13q1.1		Chr 29: 35, 752, 880–35, 896, 148	CFA29	K315 FH2385	0.15 0.03	13.45 22.89	1433 725	BH760135, BH760136; Sidjanin et al. (2002)		
<i>LHX4</i>	398-N19	13q4.4–4.6		Chr 7: 16, 740, 366–16, 782, 002	CFA7	FH2772 VIASD10	0.05 0.07	21.39 17.57	725 444	van Oost et al. (2002)		
<i>ASIP</i>	19-I2	14p1.4–1.5		Chr 24: 26, 327, 360–26, 366, 307	CFA24	CENPF REN95M13	0.14 0.14	15.70 16.21	418 653	This study	c	
<i>OTK</i> and <i>AVP</i>	15-D10	14q1.1–p1.3		OTK—Chr 24: 21, 190, 516–21, 191, 062 AVP—Chr 24: 21, 180, 172–21, 181, 841		BAC_375-K22	0.15	15.10	686	This study	n	
<i>PAX3</i>	302-P11	16p1.1–1.4		Chr 37: 31, 348, 610–31, 446, 429	CFA37 (Lowe et al. 2003)					This study		
<i>MAOA</i>	163-B12	Xp1.2–q1.1		Chr 2: 82, 718, 722–82, 809, 074 Chr X: 37, 627, 026–37, 692, 688	X	REN172J16	0.06	19.83	334	This study	n	
<i>Factor VIII (F8A)</i>	292-C4	Xq2.3–2.4		Chr X: 125, 917, 495–126, 063, 641	X	FH2997 GDI1	0.12 0.12	16.90 9.92	306 1093	Lozier et al. (2002)		
						EST16E3	0.17	9.11	1112			

The location of the same genes in the CanFam2 canine genome assembly and on the canine RH map 5000 cR is indicated. “n,” “c,” and “m” indicate genes known to be involved in neurobiology and behavior, coat color, and morphology, respectively. BAC clone 15-D10 contains at least 2 genes OTK and AVP, and BAC clone 327-P20 contains genes Tbx2 and Tbx4. LOD, logarithm of the odds.

B chromosomes (Wipf and Scackelford 1942; Gustavsson 1964; Belyaev et al. 1974). The American mink's karyotype presents 14 pairs of bi-armed chromosomes plus a pair of sex chromosomes (Mandahl and Fredga 1975; Christensen et al. 1996). The cytogenetic relationships among dog, red fox, and American mink genomes is well understood, as fluorescence in situ hybridization (FISH) with whole-chromosome probes established the correspondence of chromosomal arms between the dog and fox, and the dog and mink karyotypes (Yang et al. 1999, 2000; Graphodatsky, Yang, O'Brien et al. 2000; Graphodatsky, Yang, Serdukova et al. 2000; Graphodatsky et al. 2001, 2002).

Among the Canidae, the genome of the dog has been the most intensely studied to date. Genomic tools available for such studies include 2 bacterial artificial chromosome (BAC) libraries; meiotic, radiation hybrid (RH), and single nucleotide polymorphism maps; and an assembled genome sequence anchoring these resources (Werner et al. 1999; Breen et al. 2001, 2004; Kirkness et al. 2003; Hitte et al. 2005; Lindblad-Toh et al. 2005). The evolutionary closeness of the dog and fox provides an opportunity to apply canine genomic tools to study the fox genome. Recently, a meiotic linkage map of the silver fox (a coat color variant of the red fox) was constructed using dog-derived microsatellite markers (Kukekova et al. 2004, 2007). Alignment of the fox map against the dog genome identified highly conserved marker order between orthologous regions of dog and fox chromosomes.

Although some gene-specific markers have been placed on the fox meiotic linkage map, this approach has several limitations. Finding gene location by linkage mapping requires identification of gene-specific polymorphic markers and genotyping of these markers in reference 3-generation pedigrees. Furthermore, because alignment of the fox meiotic linkage map against the dog genome has identified pericentromeric regions of recombination suppression in fox chromosomes, marker order in these regions cannot be resolved unambiguously by meiotic linkage mapping.

Cross-species FISH mapping of BAC clones has been used previously for comparative mapping (Di Meo et al. 2005; Larkin et al. 2006; Davis et al. 2008), and several canine BAC clones have been mapped in the genome of the red fox and other canid species (Szczerebal et al. 2003, 2006; Graphodatsky et al. 2005; Klukowska-Rötzler et al. 2005). Two canine genome FISH maps are available, both anchored to the dog sequence assembly, with resolutions of 2 and 10 Mb, respectively (Thomas et al. 2005, 2007). A comparable FISH map of the fox genome would allow alignment to the dog genome at a high order of resolution than previously possible and provide a useful tool for genetic studies in foxes.

In the present study, 40 gene-containing canine BAC clones were selected to test whether such clones can be routinely used for gene mapping in foxes and were further evaluated to explore whether canine genomic tools might also be used to assign gene-specific markers in the genome of the American mink (*M. vison*), a more distantly related farm-bred species. The clones included 21 containing genes

implicated in coat color, neurobiology, and skeletal development and selected as potential candidate genes for phenotypes segregating in these species.

Materials and Methods

Selection of Canine BAC Clones for Comparative Mapping

Forty clones (Table 1) were selected from the RPCI-81 canine 8.1-fold BAC library (<http://www.chori.org/bacpac/mcanine81.htm>). Nineteen of the selected clones had been previously identified as containing specific genes (Table 1), and a further 21 clones, each positive to a single gene-specific probe, were identified specifically for this study (Table 1) by screening the library as previously described (Kukekova et al. 2003). BAC DNA was extracted by alkaline lysis (Birnboim and Doly 1979). Polymerase chain reaction (PCR) amplification of BAC DNA with gene-specific primers (Supplementary Table 1) and sequencing were used to confirm that each selected clone contained the correct corresponding gene.

Slide and Probe Preparation

Metaphase chromosomal spreads were prepared from primary fibroblast and peripheral blood lymphocyte cultures of red fox (*V. vulpes*) and American mink (*M. vison*) as described previously (Yang et al. 1999; Graphodatsky, Yang, O'Brien et al. 2000; Graphodatsky, Yang, Serdukova et al. 2000; Graphodatsky et al. 2002; Yudkin et al. 2007).

FISH Mapping

FISH on chromosomal spreads of red fox and American mink was performed using a standard protocol (Yang et al. 1999; Graphodatsky, Yang, O'Brien et al. 2000; Graphodatsky, Yang, Serdukova et al. 2000). Canine BAC clones were labeled by NICK Translation Kit (Invitrogen, Carlsbad, CA).

RH Mapping of Selected Genes on Canine 5000-centiray Panel

The 5000-centiray (cR) canine RH panel (Priat et al. 1998) was used for RH mapping of selected genes on dog chromosomes. Gene-specific primers were designed using sequences retrieved from the Trace Archive database (Supplementary Table 1). Primers for each gene were tested first on the DNA of dog and hamster, and only primer pairs that amplified the expected size product from canine but not hamster DNA were used for RH mapping. Each marker was amplified on the RH panel under the following conditions: 96 °C for 2 min; 30 cycles of 96 °C (20 s), 58 °C (20 s), and 72 °C (20 s); and a final extension at 72 °C for 5 min. All PCRs were performed using 25 ng of genomic DNA from each cell line in a final volume 15 µl, and the products were separated on a 1.8% agarose gel. PCR products were visualized by ethidium bromide staining. All markers were scored, and the complete marker set was analyzed together with 3270 markers previously mapped at




Figure 1. Examples of FISH mapping (left) of canine-derived BAC clones onto G-banded (right) metaphases of the red fox (VVU) and American mink (MVI). Localization of the TYRP2 onto VVU6p1.3–1.4 (a) and MYO15A onto MVI8p12–13 (b). Scale bar—10 μ m.

the 5000-rad panel (<http://www-recomgen.univ-rennes1.fr/Dogs/>; Breen et al. 2004) using MultiMap (Matise et al. 1994). Two-point linkage analysis was performed for each selected gene to determine its nearest markers and chromosomal location. Presence of gene-specific amplicons produced for RH mapping in the corresponding canine BAC clones was confirmed by PCR.

Localization of Selected Genes in CanFam2

To identify positions of selected genes in the 7.6 \times sequence assembly of the dog genome (CanFam2), canine orthologs were located in the canine assembly by 1) gene names and 2) RH primer sequences mapped in silico using resources of the University of Southern California, Santa Cruz, Genome Bioinformatics Site (<http://genome.ucsc.edu/>).

Results

Localization of Selected Genes in the Dog Genome

RH Mapping

Two-point linkage analysis of 38 canine genes with 3270 markers previously mapped on RH map 5000 cR identified the chromosomal location of 36 genes (Table 1). For each tested gene, the position of at least 5 RH markers that

showed linkage with the highest support was identified on the RH₅₀₀₀ map (Breen et al. 2004). Genes were assigned to a particular chromosome when all such markers mapped to the same chromosomal region. The chromosomal locations of the 2 closest markers for each mapped gene on the RH₅₀₀₀ map (Breen et al. 2004) are listed in the Table 1. RH-mapped genes were assigned to 19 dog chromosomes with 8 chromosomes containing more than one gene (Table 1). The gene *AVP* has not been mapped; all primer pairs designed for this gene produced similar size band on hamster DNA. Location of *HOXA* could not be determined unambiguously. Two genes (*PAX3* and *KDR*) have not been mapped in this study, but location of these genes on the RH₅₀₀₀ map has been identified previously (Lowe et al. 2003; Breen et al. 2004).

Localization of Selected Genes in CanFam2

The unique locations of 35 selected genes in the CanFam2 assembly were identified using gene name as a query (Table 1). Five genes (*SLC6A3*, *SLC25A3*, *ATOX1* pseudogene, *PAX3*, and *HOXA*) were assigned to more than one chromosome in CanFam2 (Table 1). Four of these genes (*SLC6A3*, *SLC25A3*, *ATOX1* pseudogene, and *PAX3*) were localized uniquely in CanFam2 using in silico PCR with primer pairs used for BAC screening or RH mapping. Location of *HOXA* could not be determined unambiguously.




Figure 2. Chromosomal locations of 40 genes mapped in the fox genome by FISH. Positions of mapped genes are indicated on the comparative chromosomal map of the red fox and dog (modified from Yang et al. 1999, 2000) to simplify comparative analysis between 2 species. Dog chromosomes are labeled based on the chromosomal nomenclature used for canine genome sequence assembly (Breen et al. 1999; Lindblad-Toh et al. 2005). Fox chromosomes are located on the left side of each panel; dog chromosomes are on the right. Positions of the genes mapped in the fox genome by FISH are indicated on the left side of fox chromosomes. Each of dog chromosomes 1, 13, 18, and 19 are syntenic to 2 fox chromosomes. Horizontal bars drawn on dog chromosomes indicate breakpoints.

For 38 genes, the canine chromosomal locations on CanFam2 and the canine RH map were in agreement (Table 1). *SLC6A3* and *AVP* were the exceptions: *SLC6A3* was assigned to CFA22 by RH mapping but is most likely located on CFA34 in CanFam2, and *AVP* was identified on CanFam2 but not on the RH Map.

Localization of Canine BAC Clones to Fox Chromosomes by FISH

All 40 canine gene-containing BAC clones were uniquely mapped to fox chromosomes by FISH. BAC clones were assigned to 15 of 17 fox chromosomes (Table 1 and Figure 1a). More than one BAC clone were assigned to 13 fox chromosomes: Two BACs were assigned to VVU1, 6, 7, 8, 9, 10, 13, 14, and X; 3 BACs were assigned to VVU3 and 12; 4 BACs were assigned to VVU4; and 10 BACs were assigned to VVU2 (Table 1 and Figure 2).

Comparative locations of 38 genes on the fox and dog chromosomes were in agreement with comparative cytogenetic data. Localization of a BAC clone containing *SLC6A3* gene on VVU2 was in disagreement with both the most

likely location of this gene in CanFam2 (CFA34) and its location predicted by RH mapping (CFA22). Because *HOXA* has not been identified in the dog genome, we could not make a comparison between dog and fox for this gene. The BAC containing the gene *C-KIT* localized to both VVU2 and B chromosomes as described previously for other *C-KIT* BAC clones (Graphodatsky et al. 2005).

These newly located genes, together with those previously mapped in the fox genome using hybrid cell lines (Rubtsov et al. 1988, 1998), increased the total number of gene-specific markers in the fox genome to 77 (Supplementary Table 2).

Localization of Canine BACs to Mink Chromosomes

Four BAC clones, containing the genes *HTR1B*, *GNGT1*, and *MYO15A* and the *ATOX1* pseudogene, were uniquely assigned to mink chromosomes (Figures 1b and 3) and in agreement with comparative cytogenetic data for mink and dog genomes (Figure 3 and Table 1). In each case, these 4 gene-containing BAC clones identified segments of mink chromosomes that were orthologous to the corresponding fox chromosomes (Figure 2, Table 1).




Figure 3. Subchromosomal localization of 4 canine BAC clones onto American mink chromosomes. Homology of American mink and dog chromosomes has been described previously (Graphodatsky, Yang, O'Brien et al. 2000; Graphodatsky, Yang, Serdukova et al. 2000; Graphodatsky et al. 2002). Numbers beside the vertical lines (left) correspond to dog chromosomes. Vertical black bars on the right side of each fox chromosome show location of canine BAC clones (this study).

Discussion

Forty canine-derived BAC clones containing known genes were uniquely assigned to fox chromosomes by FISH. Among selected BAC clones, 15 contain genes implicated in neurobiology and behavior, 5 clones contain genes involved in coat color in mammals, and at least 1 clone contains a gene important for morphology. Selection of these genes for FISH mapping in the fox genome was predetermined by the needs of ongoing genetic studies in foxes. The rest of the clones used in this study represent previously characterized BAC clones selected to increase the number of probes.

When BAC clones for FISH mapping were selected for this study, the RPCI-81 BAC library was the only available well-characterized canine BAC library. In contrast to the canine library CHORI-82 that was subsequently used for the dog genome sequencing (Lindblad-Toh et al. 2005), RPCI-81 clones are not anchored in the canine sequence assembly, and the location of each clone in the dog genome had to be identified *de novo*. To do so, both *in silico* localization of these genes in CanFam2 and RH mapping were used. The chromosomal localization of 36 of these genes in the dog was in agreement for the 2 methods. Location of 2 genes was in disagreement (Table 1). The *HOXA* gene was not uniquely localized in the dog genome by either method. The BAC clone containing *HOXA* gene had been identified by screening the RPCI-81 BAC library with the probe AF103746 (National Center for Biotechnology Information database; Li et al. 1999). This probe is 118 bp long and most likely does correspond to the *HOXA4* gene on CFA14, although the statistical support for this location is not significantly higher than for several other *HOXA* genes. RH mapping with primers based on AF103746 did not identify

a unique location for this gene in the dog genome either. For this reason, this gene was excluded from the comparative analysis of the dog and fox.

Comparison of gene locations between the fox and dog genomes was performed to test whether gene locations identified in the fox genome by FISH agreed with their chromosomal locations predicted from comparative cytogenetic data and alignment of the fox meiotic linkage map to the dog genome. The locations of 38 genes mapped in the fox genome agreed with these predictions. Location of one gene, *SLC6A3*, was in disagreement. The most likely location of this gene in CanFam2 is on CFA34, which is homologous to fox chromosome 3, but the BAC 89-K1 carrying *SLC6A3* gene was mapped to the fox chromosome 2, close to the *SLC6A4* gene. Similarity between the coding regions of these 2 genes or chimeric structure of the BAC 89-K1 can contribute to unpredicted localization of this gene. Alternatively, but less likely, the *SLC6A4* mapping results might also indicate a small insertion in VVU2 corresponding to part of CFA34 that has not yet been identified.

Previously, FISH localization of 3 canine BAC clones in 4 canids—the dog, red fox, arctic fox, and Chinese raccoon dog—has been reported (Szczerbal et al. 2003; Klukowska-Rötzler et al. 2005). These clones were selected from a different dog BAC library (<http://www.dogmap.ch/getclone.html>, Schelling et al. 2002), but 2 of them contain genes included in the current study (*IGF1* and *HTR2A*). Our results confirmed localization of the *IGF1* gene onto VVU10q chromosome and *HTR2A* on VVU6p.

In the current study, 2 genes (*TYRP2* and *HTR2A*) were mapped to the short arm of fox chromosome 6, which is homologous to CFA22. *TYRP2* mapped close to the centromere, whereas *HTR2A* mapped close to the telomere. According to the CanFam2 annotation, the location of these genes on CFA22 is opposite, *TYRP2* being near the dog telomere, whereas *HTR2A* is located close to the centromere. The relative positions of *HTR2A* and *TYRP2* genes on VVU6 indicate that the segment homologous to CFA22 has the opposite orientation in VVU6 relative to that reported by comparative chromosomal painting (Graphodatsky et al. 2001, Figure 2). This observation is also supported by previous FISH mapping results for *HTR2A* (Szczerbal et al. 2006) and by alignment of the fox meiotic linkage map against the dog genome (Kukekova et al. 2007).

Previously, more than 40 gene-specific markers have been localized in the fox genome using a panel of somatic cell hybrids (Rubtsov et al. 1988, 1998). These data, together with mapping data reported in Szczerbal et al. (2003, 2006), and FISH mapping results from the current study bring the number of genes mapped onto specific fox chromosomes to 77 (Supplementary Table 2).

In the current study, we found that FISH mapping with canine-derived BAC clones is an efficient approach for mapping genes in the silver fox genome, as all 40 canine BAC clones used in our study were uniquely assigned to fox chromosomes. Locations of 38 genes were in agreement with the cytogenetic predictions; locations of 2 genes were inconclusive. This is not unexpected as there is some

inconsistency even when canine BAC clones are FISH mapped in the dog (Breen et al. 2004). When an unpredicted location of a BAC clone is identified, additional BAC clones from the same genomic region should be mapped to understand whether observed results were caused by chimeric BAC structure, sequence similarity between different genes, or chromosomal insertions that were not detected previously. Results of this pilot project indicate an opportunity for large-scale mapping of the fox genome with canine-derived BAC clones. BAC clones from the CHORI-82 library are well characterized and anchored in the CanFam2 dog genome assembly (Lindblad-Toh et al. 2005). Mapping of the dog genome with panels of CHORI-82 BAC clones demonstrated that multicolor FISH allows construction of a gene map of the dog genome with the resolution up to 2 Mb (Thomas et al. 2005, 2007). A high-density gene map of the fox genome constructed using multicolor FISH would allow high-resolution comparison of the chromosomal regions between fox, dog, and human, refining breakpoints between homologous chromosomes, characterization of the telomeric ends of fox chromosomes, and establishing gene order between homologous regions of dog and fox chromosomes. A fox–dog comparative gene map will be an important resource for genetic studies in foxes, allowing localization of fox-mapped loci in the dog genome, selection of canine microsatellite markers for fine mapping of these loci in foxes, and identification of positional candidate genes. Furthermore, construction of high-resolution comparative FISH maps for different canid species will provide new insights into the karyotypic evolution of the Canidae.

FISH mapping results for canine-derived BAC clones in the American mink were less successful. Only 4 canine BAC clones were uniquely assigned to mink chromosomes (Figure 3, Table 1). Previous reports of FISH mapping of bovine-derived BAC clones in the genome of American mink were more encouraging (Larkin et al. 2006). These different success rates may represent differences in protocol or gene selection. A BAC library of the European ferret (*Mustela furo*) that has recently become available (<https://bacpac.chori.org/library.php?id=330>), together with the first meiotic linkage map of the American mink (Anistoroaei et al. 2007), should provide valuable resources for genetic studies in *Mustelida*. Locations of the 4 genes mapped by FISH in mink were in agreement with cytogenetic predictions based on the established homology between the dog and mink chromosomes (Yang et al. 1999; Graphodatsky, Yang, Serdukova et al. 2000; Graphodatsky et al. 2002). The fact that these 4 probes identified regions of the mink and the fox genomes, which were homologous to the same dog chromosomes, demonstrates that there is utility for this method of comparative mapping and gene identification in the 3 species.

Supplementary Material

Supplementary Tables 1 and 2 can be found at <http://www.jhered.oxfordjournals.org/>.

Funding

National Institute of Health (MH077811 to A.V.K., J.L.J., S.V.T., G.M.A.; EY006855 to G.M.A., A.V.K.; R03TW008098 to L.N.T.); Russian Fund for Basic Research to A.S.G., L.N.T., N.V.V., V.R.B., D.V.Y.; programs of the Russian Academy of Science: Molecular and Cell Biology, Biosphere Origin and Evolution, Dynamics of Genofonds of Plants, Animals and Human to A.S.G., L.N.T., N.V.V., V.R.B., D.V.Y.; Integration program of the Siberian Branch of the Russian Academy of Science to A.S.G., L.N.T., N.V.V., V.R.B., D.V.Y.

Acknowledgments

We are grateful to Dr Matthew Breen for valuable discussions on FISH mapping in canids.

References

- Andelfinger G, Etter L, Dymont M, Hitte C, Galibert F, Kirkness E, Benson DW. 2003. Radiation hybrid mapping and genomic organization of canine TBX2 and TBX4. *Anim Genet.* 34(4):307–309.
- Anistoroaei R, Menzorov A, Serov O, Farid A, Christensen K. 2007. The first linkage map of the American mink (*Mustela vison*). *Anim Genet.* 38(4):384–388.
- Belyaev DK, Trut LN, Ruvinsky AO. 1975. Genetics of the W locus in foxes and expression of its lethal effects. *J Hered.* 66(6):331–338.
- Belyaev DK, Volobuev VT, Radjabli SI, Trut LN. 1974. Polymorphism and mosaicism for additional chromosomes in silver foxes. *Genetika.* 10:58–67.
- Bininda-Emonds ORP, Gittleman J, Purvis A. 1999. Building large trees by combining phylogenetic information: a complete phylogeny of the extant *Carnivora* (*Mammalia*). *Biol Rev.* 74:143–173.
- Birnboim HC, Doly J. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7(6):1513–1523.
- Breen M, Hitte C, Lorentzen TD, Thomas R, Cadieu E, Sabacan L, Scott A, Evanno G, Parker HG, Kirkness EF, et al. 2004. An integrated 4249 marker FISH/RH map of the canine genome. *BMC Genomics.* 5(1):65–75.
- Breen M, Jouquand S, Renier C, Mellersh CS, Hitte C, Holmes NG, Chéron A, Suter N, Vignaux F, Bristow AE, et al. 2001. Chromosome-specific single-locus FISH probes allow anchorage of an 1800-marker integrated radiation-hybrid/linkage map of the domestic dog genome to all chromosomes. *Genome Res.* 11:1784–1795.
- Breen M, Thomas R, Binns MM, Carter NP, Langford CF. 1999. Reciprocal chromosome painting reveals detailed regions of conserved synteny between the karyotypes of the domestic dog (*Canis familiaris*) and human. *Genomics.* 61:145–155.
- Christensen K, Brusgaard K, Malchenko S, Lohi O, Serov O. 1996. Standardization of the American mink (*Mustela vison*) karyotype and some cosmid in situ hybridization results. *Archivos de Zootecnia.* 45:259–265.
- Davis BW, Raudsepp T, Pearks Wilkerson AJ, Agarwala R, Schäffer AA, Houck M, Ryder OA, Chowdhury BP, Murphy WJ. 2008. A high-resolution cat radiation hybrid and integrated FISH mapping resource for phylogenomic studies across Felidae. *Genomics.* 93(4):299–304.
- Debenham S, Ricketts P, Holmes NG, Thomas R, Breen M, Binns M. 2001. Physical and linkage mapping of the canine phosphate carrier (SLC25A3) and apoptotic activating factor 1 (APAF1) genes to canine chromosome 15. *Anim Genet.* 32(1):50–51.
- de Gortari MJ, Freking BA, Cuthbertson RP, Kappes SM, Keele JW, Stone RT, Leymaster KA, Dodds KG, Crawford AM, Beattie CW. 1998. A second-generation linkage map of the sheep genome. *Mamm Genome.* 9:204–209.

- Di Meo GP, Perucatti A, Floriot S, Incarnato D, Rullo R, Caputi Jambrenghi A, Ferretti L, Vonghia G, Cribiu E, Eggen A, et al. 2005. Chromosome evolution and improved cytogenetic maps of the Y chromosome in cattle, zebu, river buffalo, sheep and goat. *Chromosome Res.* 13(4):349–355.
- Drögemüller C, Rak SG, Kuiper H, Leeb T, Quignon P, Galibert F, Distl O. 2002. Assignment of the canine tectorin alpha gene (TECTA) to CFA5q12–>q13 by FISH and confirmation by radiation hybrid mapping. *Cytogenet Genome Res.* 97(1–2):140A.
- Georges M. 2007. Mapping, fine mapping, and molecular dissection of quantitative trait loci in domestic animals. *Annu Rev Genomics Hum Genet.* 8:131–162.
- Graphodatsky AS, Kukekova AV, Yudkin DV, Trifonov VT, Vorobieva NV, Beklemisheva VR, Perelman PL, Graphodatskaya DA, Trut LN, Yang F, et al. 2005. The proto-oncogene C-KIT maps to canid B-chromosomes. *Chromosome Res.* 13:113–122.
- Graphodatsky AS, Yang F, O'Brien PC, Perelman P, Milne BS, Serdukova N, Kawada SI, Ferguson-Smith MA. 2001. Phylogenetic implications of the 38 putative ancestral chromosome segments for four canid species. *Cytogenet Cell Genet.* 92:243–247.
- Graphodatsky AS, Yang F, O'Brien PC, Serdukova N, Milne BS, Trifonov V, Ferguson-Smith MA. 2000. A comparative chromosome map of the Arctic fox, red fox and dog defined by chromosome painting and high resolution G-banding. *Chromosome Res.* 8:253–263.
- Graphodatsky AS, Yang F, Perelman PL, O'Brien PCM, Serdukova NA, Milne BS, Biltueva LS, Fu B, Vorobieva NV, Kawada SI, et al. 2002. Comparative molecular cytogenetic studies in the order Carnivora: mapping chromosomal rearrangements onto the phylogenetic tree. *Cytogenet Genome Res.* 96:137–145.
- Graphodatsky AS, Yang F, Serdukova N, Perelman P, Zhdanova N, Ferguson-Smith MA. 2000. Dog chromosome-specific paints reveal evolutionary inter- and intrachromosomal rearrangements in the American mink and human. *Cytogenet Cell Genet.* 90:275–278.
- Gustavsson I. 1964. The chromosomes of the dog. *Hereditas.* 51:187–189.
- Haworth K, Putt W, Cattanaach B, Breen M, Binns M, Lingaas F, Edwards YH. 2001. Canine homolog of the T-box transcription factor T; failure of the protein to bind to its DNA target leads to a short-tail phenotype. *Mamm Genome.* 12(3):212–218.
- Hitte C, Madeoy J, Kirkness EF, Priat C, Lorentzen TD, Senger F, Thomas D, Derrien T, Ramirez C, Scott C, et al. 2005. Facilitating genome navigation: survey sequencing and dense radiation-hybrid gene mapping. *Nat Rev Genet.* 6:643–648.
- Karlsson EK, Baranowska I, Wade CM, Salmon Hillbertz NH, Zody MC, Anderson N, Biagi TM, Patterson N, Pielberg GR, Kulbokas EJ 3rd, et al. 2007. Efficient mapping of mendelian traits in dogs through genome-wide association. *Nat Genet.* 39(11):1321–1328.
- Kharlamova AV, Faleev VI, Trapezev OV. 2000. Effect of selection for behavior on the cranial traits of the American mink (*Mustela vison*). *Genetika.* 36(6):823–828.
- Kharlamova AV, Trut LN, Carrier DR, Chase K, Lark KG. 2007. Genetic regulation of canine skeletal traits: trade-offs between the hind limbs and forelimbs in the fox and dog. *Integr Comp Biol.* 47(3):373–381.
- Kirkness EF, Bafna V, Halpern AL, Levy S, Remington K, Rusch DB, Delcher AL, Pop M, Wang W, Fraser CM, et al. 2003. The dog genome: survey sequencing and comparative analysis. *Science.* 301:1898–1903.
- Klukowska-Rötzler J, Szczerbal I, Braunschweig M, Switonski M, Schelling C, Dolf G. 2005. Mapping and development of four microsatellite markers for the canine 5'-hydroxytryptamine serotonin receptor 2A (HTR2A). *Anim Genet.* 36(2):173–175.
- Kuiper H, Rak SG, Drögemüller C, Leeb T, Quignon P, Galibert F, Distl O. 2002. Assignment of the canine cadherin related 23 gene (CDH23) to chromosome 4q12–>q13 by fluorescence in situ hybridization and radiation hybrid mapping. *Cytogenet Genome Res.* 97(1–2):140B.
- Kukekova AV, Aguirre GD, Acland GM. 2003. Cloning and characterization of canine SHARP1 and its evaluation as a positional candidate for canine early retinal degeneration (erd). *Gene.* 312:335–343.
- Kukekova AV, Trut LN, Oskina IN, Johnson JL, Temnykh SV, Kharlamova AV, Shepeleva DV, Gulievich RG, Shikhevich SG, Graphodatsky AS, et al. 2007. A meiotic linkage map of the silver fox, aligned and compared to the canine genome. *Genome Res.* 17(3):387–399.
- Kukekova AV, Trut LN, Oskina IN, Kharlamova AV, Shikhevich SG, Kirkness EF, Aguirre GD, Acland GM. 2004. A marker set for construction of a genetic map of the silver fox (*Vulpes vulpes*). *J Hered.* 95:185–194.
- Larkin DM, Prokhorovich MA, Astakhova NM, Zhdanova NS. 2006. Comparative mapping of mink chromosome 8p: in situ hybridization of seven cattle BAC clones. *Anim Genet.* 37(4):429–430.
- Li R, Mignot E, Faraco J, Kadotani H, Cantanese J, Zhao B, Lin X, Hinton L, Ostrander EA, Patterson DF, et al. 1999. Construction and characterization of an eightfold redundant dog genomic bacterial artificial chromosome library. *Genomics.* 58(1):9–17.
- Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, Clamp M, Chang JL, Kulbokas EJ 3rd, Zody MC, et al. 2005. Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature.* 438:803–819.
- Lowe JK, Kukekova AV, Kirkness EF, Langlois MC, Aguirre GD, Acland GM, Ostrander EA. 2003. Linkage mapping of the primary disease locus for collie eye anomaly. *Genomics.* 82(1):86–95.
- Lozier JN, Dutra A, Pak E, Zhou N, Zheng Z, Nichols TC, Bellinger DA, Read M, Morgan RA. 2002. The Chapel Hill hemophilia A dog colony exhibits a factor VIII gene inversion. *Proc Natl Acad Sci USA.* 99(20):12991–12996.
- Matise TC, Perlin M, Chakravarti A. 1994. Automated construction of genetic linkage maps using an expert system (MultiMap): a human genome linkage map. *Nat Genet.* 6:384–390.
- Mandahl N, Fredga K. 1975. Q, G, and C-band patterns of mink chromosomes. *Hereditas.* 81:211–220.
- Nes NN, Einarsson EJ, Lohi O, Jorgensen G. 1988. Beautiful fur animals—and their color genetics. Hillerød (Denmark): SCIENTIFUR. p. 271.
- Pontius JU, Mullikin JC, Smith DR, Agencourt Sequencing Team, Lindblad-Toh K, Gnerre S, Clamp M, Chang J, Stephens R, Neelam B, et al. 2007. Initial sequence and comparative analysis of the cat genome. *Genome Res.* 17(11):1675–1689.
- Priat C, Hitte C, Vignaux F, Renier C, Jiang Z, Jouquand S, Chéron A, André C, Galibert F. 1998. A whole-genome radiation hybrid map of the dog genome. *Genomics.* 54(3):361–378.
- Rak SG, Drogemüller C, Kuiper H, Leeb T, Quignon P, Andre C, Distl O. 2002. Cloning and chromosomal localization of MYO15A to chromosome 5 of the dog (*Canis familiaris*). *Chromosome Res.* 10(5):407–410.
- Reed KM, Chaves LD, Hall MK, Knutson TP, Harry DE. 2005. A comparative genetic map of the turkey genome. *Cytogenet Genome Res.* 111:118–127.
- Robinson R. 1975. The American mink (*Mustela vison* Schreber). In: Robinson R, editor. *Handbook of genetics* 4. New York: Plenum. p. 367–398.
- Rogers J, Garcia R, Shelledy W, Kaplan J, Arya A, Johnson Z, Bergstrom M, Novakowski L, Nair P, Vinson A, et al. 2006. An initial genetic linkage map of the rhesus macaque (*Macaca mulatta*) genome using human microsatellite loci. *Genomics.* 87:30–38.
- Rogers J, Mahaney MC, Witte SM, Nair S, Newman D, Wedel S, Rodriguez LA, Rice KS, Slifer SH, Perelygin A, et al. 2000. A genetic linkage map of the baboon (*Papio hamadryas*) genome based on human microsatellite polymorphisms. *Genomics.* 67:237–247.
- Rubtsov N, Graphodatsky A, Matveeva VG, Radjabli SI, Nesterova TB, Kulbakina NA, Zakian S. 1988. Silver fox gene mapping: conserved

- chromosome regions in the order *Carnivora*. *Cytogenet Cell Genet*. 48:95–98.
- Rubtsov NB. 1998. The fox gene map. *ILAR J*. 39:92–95.
- Shackelford RM. 1948. The nature of coat color differences in mink and foxes. *Genetics*. 33:311–336.
- Shackelford RM. 1949. Six mutations affecting coat color in ranch bred mink. *Am Nat*. 83:49–86.
- Schelling C, Schl pfer J, Billault A, Guziewicz K, Gm r A, Katmann I, Pineroli B, Colomb B, Rickli O, Wittwer C, et al. 2002. Construction of a canine bacterial artificial chromosome library for screening with PCR. *J Anim Breed Genet*. 119:400–401.
- Sidjanin DJ, Lowe JK, McElwee JL, Milne BS, Phippen TM, Sargan DR, Aguirre GD, Acland GM, Ostrander EA. 2002. Canine CNGB3 mutations establish cone degeneration as orthologous to the human achromatopsia locus ACHM3. *Hum Mol Genet*. 11(16):1823–1833.
- Sidjanin DJ, Miller B, Kijas J, McElwee J, Pillardy J, Malek J, Pai G, Feldblyum T, Fraser C, Acland G, et al. 2003. Radiation hybrid map, physical map, and low-pass genomic sequence of the canine prcd region on CFA9 and comparative mapping with the syntenic region on human chromosome 17. *Genomics*. 81(2):138–148.
- Szczerbal I, Klukowska-Roetzler J, Dolf G, Schelling C, Switonski M. 2006. FISH mapping of 10 canine BAC clones harbouring genes and microsatellites in the arctic fox and the Chinese raccoon dog genomes. *J Anim Breed Genet*. 123(5):337–342.
- Szczerbal I, Rogalska-Niznik N, Klukowska J, Schelling C, Dolf G, Switonski M. 2003. Comparative chromosomal localization of the canine-derived BAC clones containing LEP and IGF1 genes in four species of the family *Canidae*. *Cytogenet Genome Res*. 102:264–266.
- Thomas R, Duke SE, Bloom SK, Breen TE, Young AC, Feiste E, Seiser EL, Tsai PC, Langford CF, Ellis P, et al. 2007. A cytogenetically characterized, genome-anchored 10-Mb BAC set and CGH array for the domestic dog. *J Hered*. 98(5):474–484.
- Thomas R, Scott A, Langford CF, Fosmire SP, Jubala CM, Lorentzen TD, Hitte C, Karlsson EK, Kirkness E, Ostrander EA, et al. 2005. Construction of a 2-Mb resolution BAC microarray for CGH analysis of canine tumors. *Genome Res*. 15(12):1831–1837.
- Trapezov OV, Trapezova LI, Sergeev EG. 2008. Effect of coat color mutations on behavioral polymorphism in farm populations of American minks (*Mustela vison* Schreber, 1777) and sables (*Martes zibellina* Linnaeus, 1758). *Genetika*. 44(4):516–523.
- Trut LN. 1999. Early canid domestication: the farm fox experiment. *Am Sci*. 87:160–169.
- Trut LN. 2001. Experimental studies of early canid domestication. The genetics of the dog. Oxford: CABI. p. 15–43.
- V ge DI, Lu D, Klungland H, Lien S, Adalsteinsson S, Cone RD. 1997. A non-epistatic interaction of agouti and extension in the fox, *Vulpes vulpes*. *Nat Genet*. 15(3):311–315.
- van de Sluis B, Nanji MS, Breen M, Pearson PL, Oost BA, Cox DW, Wijnga C. 2001. Characterization and chromosomal localization of five canine ATOX1 pseudogenes. *Cytogenet Cell Genet*. 93(1–2):105–108.
- van den Berg L, Imholz S, Versteeg SA, Leegwater PA, Zijlstra C, Bosma AA, van Oost BA. 2004. Isolation and characterization of the canine serotonin receptor 1B gene (htr1B). *Gene*. 326:131–139.
- van Oost BA, Versteeg SA, Imholz S, Kooistra HS. 2002. Exclusion of the lim homeodomain gene LHX4 as a candidate gene for pituitary dwarfism in German shepherd dogs. *Mol Cell Endocrinol*. 197(1–2):57–62.
- Wayne RK, Geffen E, Girman DJ, Koepfli KP, Lau LM, Marshall CR. 1997. Molecular systematics of the Canidae. *Syst Biol*. 46:622–653.
- Wayne RK, Ostrander EA. 2007. Lessons learned from the dog genome. *Trends Genet*. 23(11):557–567.
- Werner P, Mellersh CS, Raducha MG, DeRose S, Acland GM, Prociuk U, Wiegand N, Aguirre GD, Henthorn PS, Patterson DF, et al. 1999. Anchoring of canine linkage groups with chromosome-specific markers. *Mamm Genome*. 10:814–823.
- Wipf L, Scackelford RM. 1942. Chromosomes of red fox. *Proc Natl Acad Sci USA*. 28:265–268.
- Yang F, Milne BS, Schelling C, Dolf G, Schl pfer J, Switonski M, Ladon D, Pienkowska A, Bosma AA, Sargan DR, et al. 2000. Chromosome identification and assignment of DNA clones in the dog using a red fox and dog comparative map. *Chromosome Res*. 8:93–100.
- Yang F, O'Brien PC, Milne BS, Graphodatsky AS, Solanky N, Trifonov V, Rens W, Sargan D, Ferguson-Smith MA. 1999. A complete comparative chromosome map for the dog, red fox, and human and its integration with canine genetic maps. *Genomics*. 62:189–202.
- Yudkin DV, Trifonov VA, Kukekova AV, Vorobieva NV, Rubtsova NV, Yang F, Acland GM, Ferguson-Smith MA, Graphodatsky AS. 2007. Mapping of KIT adjacent sequences on canid autosomes and B chromosomes. *Cytogenet Genome Res*. 116(1–2):100–103.

Received November 22, 2008; Revised March 24, 2009;
Accepted May 13, 2009

Corresponding Editor: Dr. Elaine Ostrander